

# Staff Nurse Model Question Paper

---

**1. For education of measles, immunization coverage should be at least**

- (a) 80%
- (b) 85%
- (c) 96%
- (d) 100%

**Answer : B**

**2. Period of maximum infectivity in mumps is during**

- (a) incubation period
- (b) onset of swelling
- (c) after swelling appears
- (d) prodromal period

**Answer : A**

**3. Major epidemics of influenza A occur at intervals of**

- (a) 1 year
- (b) 2-3 years
- (c) 10-15 years
- (d) 20-25 years

**Answer : B**

**4. The most common source of infection for diphtheria is a**

- (a) case
- (b) sub clinical case
- (c) carrier
- (d) all of the above

**Answer : D**

**5. Shick test is done in**

- (a) meningitis
- (b) diphtheria
- (c) pertussis
- (d) polimyelitis

**Answer : A**

**6. Adjuvant present in DPT vaccine is**

- (a) aluminium phosphate
- (b) aluminium hydroxide
- (c) zinc phosphate
- (d) magnesium phosphate

**Answer : C**

**7. Recommended site of administration of DPT in infants is**

- (a) gluteal
- (b) deltoid
- (c) lateral aspect of thigh
- (d) forearm

**Answer : C**

**8. The most severe complications following DPT are due to ..... Component**

- (a) diphtheria
- (b) pertussis
- (c) tetanus
- (d) adjuvant

**Answer: D**

**9. The vaccine usually given to pilgrims to the Middle East is**

- (a) DPT
- (b) influenza
- (c) yellow fever
- (d) meningitis

**Answer : C**

**10. Best index to estimate case load of T.B. in a community is**

- (a) incidence of infection
- (b) prevalence of infection
- (c) incidence of disease
- (d) prevalence of disease

**Answer : C**

**11. Prevalence of T.B. infection in a community is estimated by**

- (a) tuberculin test
- (b) sputum microscopy
- (c) culture
- (d) chest X-ray

**Answer : A**

**12. The amount of PPD used in mantoux test in India is**

- (a) 1 TU
- (b) 5 TU
- (c) 25 TU
- (d) 250 TU

**Answer : A**

**13. Result of mantoux test is read after**

- (a) 6 hours
- (b) 24 hours
- (c) 48 hours
- (d) 72 hours

**Answer : C**

**14. Present day BCG vaccine is supplied as**

- (a) liquid vaccine
- (b) freeze dried vaccine
- (c) both
- (d) none of the above

**Answer : B**

**15. The diluent used for reconstituting BCG vaccine is**

- (a) normal saline
- (b) distilled water
- (c) dextrose
- (d) none of the above

**Answer : A**

**16. Reconstituted BCG vaccine should be used within**

- (a) 3 hours
- (b) 6 hours
- (c) 18 hours
- (d) 24 hours