DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

Test Booklet Series

TEST BOOKLET

T. B. C.: HM - 1/19

1753



HOMOEOPATHIC MEDICAL OFFICER SI. No.

PAPER -- I

Time Allowed: 12 Hours

Maximum Marks: 100

- : INSTRUCTIONS TO CANDIDATES : IMMEDIA TELY AFTER THE COMMENCEMENT OF THE EXAMINATION, YOU SHOULD CHECK THAT 1. THIS TEST BOOKLET DOES NOT HAVE ANY UNPRINTED OR TORN OR MISSING PAGES ORITEMS ETC. 1 IF SO, GETIT REPLACED BYACOMPLIETE TEST BOOKLET OF THE SAME SERIES ISSUED TO YOU. 2. ENCODE CLEARLY THE TEST BOOKLET SERIES A, B, C OR D, AS THE CASE MAY BE, IN THE APPROPRIATE PLACE IN THE ANSWER SHEET USING BALL POINT PEN (BLUE OR BLACK). 3 You have to enter your Roll No. on the Test Booklet in the Box provided alongside. DO NOT write anything else on the Test Booklet. 4 YOU ARE REQUIRED TO FILL UP & DARKEN ROLL NO., TEST BOOKLET / QUESTION BOOKLET SERIES IN THE ANSWER SHEET AS WELL AS FILL UP TEST BOOKLET / QUESTION BOOKLET SERIES AND SERIAL NO. AND ANSWER SHEET SERIAL NO. IN THE A TTENDANCE SHEET CAREFULLY. WRONGLY FILLED UP ANSWER SHEETS ARE LIABLE FOR **REJECTION AT THE RISK OF THE CANDIDATE.** This Test Booklet contains 100 items (questions). Each item (question) comprises four 5. responses (answers). You have to select the correct response (answer) which you want to mark (darken) on the Answer Sheet. In case, you feel that there is more than one correct response (answer), you should mark (darken) the response (answer) which you consider the best. In any case, choose ONLY ONE response (answer) for each item (question). You have to mark (darken) all your responses (answers) ONLY on the separate Answer 6 Sheet provided by using BALL POINT PEN (BLUE OR BLACK). See instructions in the Answer Sheet. All items (questions) carry equal marks. All items (questions) are compulsory. Your total 7. marks will depend only on the number of correct responses (answers) marked by you in the Answer Sheet. Before you proceed to mark (darken) in the Answer Sheet the responses to various items 8. (questions) in the Test Booklet, you have to fill in some particulars in the Answer Sheet as per the instructions sent to you with your Admission Certificate. After you have completed filling in all your responses (answers) on the Answer Sheet and after 9. conclusion of the examination, you should hand over to the Invigilator the Answer Sheet issued to you. You are allowed to take with you the candidate's copy / second page of the Answer Sheet along with the Test Booklet, after completion of the examination, for your reference.
 - 10. Sheets for rough work are appended in the Test Booklet at the end.

DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

1. Ptosis is caused by the paralysis of: III cranial nerve (A) IV cranial nerve (B) 6. V cranial nerve **(C)** VI cranial nerve (D) Acute complication of diabetes is: 2. (A) Coma **(B)** Blindness 7. Diabetes Ketoacidosis & Coma (C)(D) Heart failure Egg-shell calcification of hilar lymph 3 nodes is characteristically seen in: Tuberculosis {A} Sarcoidosis (8) 8. (C) Tieratoma (D) Silicosis Best artery for coronary angiography : 4 (A) Brachial 9. (B)Femioral Carotid (C)(D) Axillary The encephalopathy is commonly 5.seen in chronic alcoholics: (A) Reye's syndrome Multicystic enc a phalopathy (8) CC- 1A/14 (2)

Wernike's encephalopathy (C) Spongiform encephalopathy (D) All of the following may be associated with rheumatic fever except : (A) Nephritis **(B)** Chorea Polyarthritis (C) (D) Carditi 8 Haemoptysis with renal involvement in seen in: (A) Good Pasture's Syndrome Rheumatchid Avithmis (8)(C) **Reiter's Syndrome Behcet's Disease** (D) Maize eating is related to : (A) Pellagra Scurvy. (8)Beri beri (C) (D) Phrynoderma The outstanding symptom of acute pancreatitis : (A) Abdominal Bloating Severe constant upper (B) abdominal pain radiates to back Jaundice (C)Constipation (D)

Contd

· · i

- 10. The most serious complication of herpes simplex virus is :
 - (A) Leukemia
 - (B) Encephalitis
 - (C) HIV
 - (D) Rubella
- 11. The following are true regarding Lyme disease except:
 - (A) It is transmitted by Ixodid ticks
 - (B) Erythema migrans
 - (C) Borrelia recurrent is the etiological agent
 - (D) Borrelia burgdorferii is the etiological agent
- 12. Rheumatoid arthritis associated with pneumoconiosis is known as :
 - (A) Middle lobe syndrome
 - (B) Black lung syndrome
 - (C) Caplans syndrome
 - (D) Hamannrich syndrome
- 13. Ankylosing spondylitis is characterised by all except :
 - (A) Possible fusion of entire spine

CC - 1A/14

- (B) Involvement of sacroiliac joints
- (C) High prevalence of HLA-B27
- (D) Highest frequency of older women
- 14. Torres inclusion body may be found in:
 - (A) Yellow fever
 - (B) Dengue
 - (C) Kyasanur forest disease
 - (D) All of these
- 15. The initial complaint in primary biliary cirrhosis is :
 - (A) Pain in abdomen
 - (B) Fever
 - (C) Weakness
 - (D) Pruritus
- 16. Black water fever is:
 - (A) Related to heavy parasitaemia
 - (B) Always associated with renal failure
 - (C) Related to Haemoglobinuria
 - (D) Observed in incompatible blood transfusion

(Turn over)

(3)

17. Hereditary haemochromatosis is associated with :

(A) HLA-A locus on chromosome 6p

والمعالم والمعارين

- (B) HLA-B8
- (C) HLA-B7
- (D) HLA-B27
- 18. The parathyroid glands are:
 - (A) 3
 - (B) 2
 - (C) 4
 - (D) 6
- 19. Pernicious anaemia is associated with:
 - (A) Hypogammaglobulinaemia
 - (B) Blood group B
 - (C) HLA-B27
 - (D) Yellow sclera
- 20. The characteristic feature of hypertrophic osteoarthropathy is :
 - (A) Clubbing
 - (B) Pallor
 - (C) Muddy sclera
 - (D) Anaemia
- CC- 1A/14

21. Streak ovary is associated with :(A) Klinefelter's Syndrome

- (B) Asherman's Syndrome
- (C) Turner's Syndrome
- (D) Super Female
- 22. Haematometra seen in association with:
 - (A) Cryptomenorrhoea
 - (B) Congestive Dysmenorrhoea
 - (C) Oligomenorrhea
 - (D) D.U.B.
- 23. Powderburn's seen in :
 - (A) Tuberculous endosalpingiosis
 - (B) Pelvic endometriosis
 - (C) Tuberculous cervicitis
 - (D) Adenomyosis
- 24. Basal body temperature noted in:
 - (A) Ovulation
 - (B) Pregnancy
 - (C) Vaginal infection
 - (D) Menopause

(4)

Contd.

| 25. Fixed retroversion seen in | ו | ; |
|--------------------------------|---|---|
|--------------------------------|---|---|

- (A) PID
- (B) Uterine Prolapse
- (C) Adenomyosis
- (D) Fibroid uterus
- 26. Culdocentesis is a method of diagnosis in :
 - (A) Fibroiduterus
 - (B) Adenomyosis
 - (C) Ovarian Tumour
 - (D) Ectopic gestation
- 27 Complete mole associated with :
 - (A) Hydatidiform mole
 - (B) Choriola
 - (C) Missed abortion
 - (D) None of these
- 28. Mastitis followed by breast abscess is most frequently due to:
 - (A) Bacterial Vaginosis
 - (B) Pneumococcus
 - (C) E coli
 - (D) Staphylococcus
- CC- 1A/14

- 29. One of them is not a complication of Placenta Previa:
 - (A) Malpresentation
 - (B) Postmaturity
 - (C) Preterm labour
 - (D) Intrapartum Haemorrhage
- 30. Which sign is absent in abdominal pregnancy?
 - (A) Braxton Hick's sign
 - (B) Foetal movements
 - (C) Foetal heart sounds
 - (D) None of these
- 31. Commonest site of Branchial Cyst
 - (A) Lower 1/3 of sternomastoid on anterior border
 - (B) Lower 1/3 of sternomastoid on posterior border
 - (C) Upper 1/3 of sternomastoid on anterior border
 - (D) Upper 1/3 of sternomastoid on posterior border
 - (Turn over)

(5)

| 32. | Most common coagulopathy noted | in | (C) Rectum managements |
|-----|------------------------------------|-----|--|
| | surgical patient is : | | (D) Colon |
| | (A) Thrombocytopaenia | 36. | Which of the following gas is a |
| | (B) Afibrinogenemia | | radiosensitiser? |
| | (C) Fibrinolysis | | (A) CO ₂ |
| | (D) Factor VIII deficiency | · , | (B) N_2 |
| 33. | Most important step in Cardia | IC. | (C) O ₂ |
| | resuscitation is : | | A state of the sta |
| | (A) Bicarbonate administration | | (D) CO |
| | (B) Cardiac massage | 37. | In obstruction of large gut, rapture |
| | (C) Ventilation | | occurs at the: |
| | (D) Intracardiac adrenalin | | (A) Caecum |
| 34. | Hour glass stomach is seen in: | ÷ | (B) Ascending colon |
| | (A) Gastric carcinoma | | (C) Transverse colon |
| | (B) Gastric ulcer | | (D) Descending colon |
| | (C) Gastric lymphoma | 38. | Most common post operative |
| ••• | (D) Corrosive stricture | | complication of Spleenectomy is: |
| 35. | Commonest site for carcinoi d tumo | ur | (A) Left side pleural effusion |
| | is: | | (B) Subphrenic abscess |
| | (A) Ileum | | (C) Left lower lobe atelectasis |
| | (B) Appendix | | (D) Pancreatic fistula |
| CC- | - 1A/14 | (6) | Contd |

- 39. Which is the commonest tumour of urinary bladder?
 - (A) Papilloma
 - (B) Adenocarcinoma
 - (C) Transitional cell carcinoma
 - (D) Squamous cell carcinoma
- 40. "Rasberry tumour" is another name for:
 - (A) Umbilical fistula
 - (B) Umbilical granuloma
 - (C) Umbilical adenoma
 - (D) Meckel's diverticulum
- 41. Commonest cause of deafness in children under 12 years :
 - (A) Trauma
 - (B) Wax
 - (C) Otosclerosis
 - (D) Secretory otitis media
- 42. CSF Rhinorrhoea is usually due to fracture of:
 - (A) Ethmoidal bone
 - (B) Cribriform plate
 - (C) Orbital plate
 - (D) Maxilla

- 43. Constantly changing refractory error is seen in:
 - (A) Traumatic cataract
 - (B) Diabetic cataract
 - (C) Morgagnian cataract
 - (D) Intumescent cataract
- 44. Commonest tumour arising from the metaphysis is :
 - (A) Osteoclastoma
 - (B) Ewing's sarcoma
 - (C) Osteosarcoma
 - (D) Synovial sarcoma
- 45. In rheumatoid arthritis, the part which is affected most is :
 - (A) Synovium
 - (B) Subchondral bone
 - (C) Cartilage
 - (D) Tendon
- 46. The concept "Levels of Prevention" in the health care systems aims at:
 - (A) Identification of stages of a disease
 - (B) Control and eradication of a disease the state section of a
 - (C) Preventing the endemic and epidemic risks
 - (D) Intervention in the natural
 - history of the disease

(7)

(Turn over)

CC- 1A/14

- 47. All are true of Soyabean execept:
 - (A) 40% Protein + 20% Fat
 - (B) Lysine deficient
 - (C) Protein of high nutritive value
 - (D) 4% mineral
- 48. All are true about attack rate except:
 - (A) Is a type of prevalence rate
 - (B) Expressed as percentage
 - (C) Used when risk period is short
 - (D) Reflects extent of epidemic
- 49. Following are true regarding chlorination except:
 - (A) Contact period -30 minutes
 - (B) Water should not be turbid
 - (C) Residual chlorine 0.5 mg/L.
 - (D) Chlorine demand should be estimated
- 50. Which of the following is not true of Cohort Study ?
 - (A) Expensive
 - (B) Incidence can be calculated
- CC- 1A/14

- (C) Always prospective
- (D) Used to study chronic disease
- 51. Which of the following is correct regarding "Health for All by 2000 AD" goals?
 - (A) NRR=1
 - (B) Crude Death Rate 9/1000 LB
 - (C) To provide potable water to rural population
 - (D) All of these
- 52. All the following are true about surveillance except:
 - (A) It is a one time observation
 - (B) Continuous scrutiny of occurrence of disease
 - (C) Essential for prevention and control
 - (D) Provides information on changing needs
- 53. Strength of sewage is expressed in terms of:
 - (A) Biological of oxygen demand
 - (B) Suspended solids
 - (C) Chemical oxygen demand
 - (D) E coli count

(8)

Contd.

- 54. Sensitivity of a test is given by :
 - (A) True positive / True positive +False positive × 100
 - (B) True positive / True positive +False negative
 - (C) True negative / False positive+ True negative
 - (D) False positive / False positive+ True negative × 100
- 55. The 'Clean 5' principles of safe motherhood includes all except :
 - (A) Clean tie
 - (B) Clean cord
 - (C) Clean perineum
 - (D) Clean room
- 56. Health Education :
 - (A) Develops reflexible behaviour
 - (B) Trains people to use judgementfor health action
 - (C) Appeals to emotions to adopt health measures
 - (D) Does not change attitude
- CC- 1A/14

- 57. All are true regarding measles except:
 - (A) Sec. attack rate 30-40%
 - (B) Carrier state not present
 - (C) Severe in malnourished child
 - (D) I attack confers life long immunity
- 58. Carriers are not source in the transmission of:
 - (A) Poliomyelitis
 - (B) Typhoid
 - (C) Diphtheria
 - (D) Measles
- 59. All of the following are considered high risk factors for development of protein energy malnutrition except :
 - (A) Measles
 - (B) Acute diarrhoea
 - (C) Tuberculosis
 - (D) Low birth weight
- 60. Keta themometer is used now-a-days to determine :
 - (A) Direction of air flow
 - (B) Air velocity
 - (C) Cooling power of air
 - (D) Humidity of air

(Turn over)

(9)

61. One of the following is peripheral nerve poison :

- (A) Conjum
- (B) Digitalis
- (C) Oleander
- (D) Aconite
- 62. Among the following irritant poison one is non-metallic poison :
 - (A) Arsenic
 - (B) Lead
 - (C) Mercury
 - (D) Phosphorus
- 63. Rule of nine estimate percentage of burns by:
 - (A) Body surface area percentage by age
 - (B) Cause of burn
 - (C) Age of person
 - (D) Surface area of burns
- 64. One of the following is typical drowning :
 - (A) Dry drowning

- (B) Immersion syndrome
- (C) Submersion of the unconsciousness
- (D) Wet drowning
- 65. In dead born child following sign may be absent:
 - (A) Scalp oedema
 - (B) Rigor mortis
 - (C) Maceration
 - (D) Mummification
- 66. For disputed paternity and maternity' following systems are used preferentially:
 - (a) ABO grouping
 - (b) MN system
 - (c) Rh sub-grouping

 - (d) Salivary DNA study
 - (A) (a), (b), (c), (d)
 - (B) (a), (c), (d), (b)
 - (C) (c), (a), (d), (b)
 - (D): (b), (c), (d), (a)

(10)

CC-1A/14

Contd

67. For certification of mental illness patient is observed for at a time :

(A) For 10 to 30 days

- (B) For 4 days
- (C) For 9 days
- (D) For 30 days

68. One of following is criminal cause of death in new born :

- (A) Immaturity
- (B) Malformation
- (C) Diseases of mother and child
- (D) Act of commission or omission
- 69. Authorisation for a medico legal autopsy is given by:
 - (A) Coroner
 - (B) Police
 - (C) Magistrate
 - (D) All of these
- 70. The presence of four wisdom teeth usually means that the subject has passed the age of:
 - (A) 10 years
 - (B) 12 years
 - (C) 17 years
 - (D) 25 years

CC- 1Å/14

71. Serotonin is produced by:

- (A) Mastcell
- (B) Macrophage
- (C) Platelet
- (D) All of these

72. Sudan IV and Oil Red O are the common stains used in :

- (A) Amyloidosis
- (B) Fatty Liver
- (C) Leprosy
- (D) All of these

73. Heart failure cell is a characteristic feature of:

- (A) Acute Pulmonary Congestion
- (B) Chronic Passive Congestion of Lung
- (C) Myocardial Infraction
- (D) Myxoma
- 74. Most Common chromosomal disorder cause mental disorder with microcephaly:
 - (A) Down syndrome
 - (B) Edward syndrome
 - (C) Felty syndrome
 - (D) All of these

(11)

(Turn over)

| dissemination of Carcinona is: (D) Normal structure of Bone marrow (A) Seeding of body cavitied 80. Codman Triangle is a feature of: (B) Implantation (A) Osteosarcoma (C) Haematogenous (B) Osteochondroma (D) Lymphatic pathway (C) Chondrosarcoma 76. Pulseless disease is: (D) Goitre (A) PAN 81. The sacro-liac joint transmit from the lower limbs to the : (D) Raynauds phenomena (A) Sacro-liac ligament (D) Raynauds phenomena (C) Lumbosacral ligament (D) Raynauds phenomena (D) Vertebral column of: 82: Lateral boundary of cubital fossa is formed by: (B) Rheumatic fever (D) Pronator teres (D) Peptic Ulcer (C) Brachialis 78. Reid Index is related to : (A) Bronchial Asthma 83. (B) Lipoma (A) Lat. half by median, i.at. half by uhar (D) Bronchitis (B) Medial half by median nerve (B) Rables (D) Whole by uhar nerve | 75. C | common pathway for initi | al | , - | (C) Endocarditis |
|---|--------------|---------------------------------------|----|-----|---|
| (B) Implantation (C) Haematogenous (D) Lymphatic pathway (C) Chondrosarcoma (D) Lymphatic pathway (C) Chondrosarcoma (D) Goitre 81. The sacro-Iliac joint transmit from the lower limbs to the : (A) PAN (B) Burger's disease (C) Takayasus disease (D) Raynauds phenomena (C) Takayasus disease (D) Raynauds phenomena (C) Lumbosacral ligament (D) Vertebral column (D) Vertebral column (E) Peptic Ulcer (C) Cirrhosis of liver (D) Peptic Ulcer (C) Brachialis (D) Peptic Ulcer (C) Leukemia (D) Bronchitis (P) Bronchitis (P) Mallory bodies are seen in : (A) Alcoholic Cirrhosis (C) Whole by median nerve | d | issemination of Carcinona is: | | | (D) Normal structure of Bone marrow |
| (B) Implantation (A) Osteosarcoma (C) Haematogenous (B) Osteochondroma (D) Lymphatic pathway (C) Chondrosarcoma 76. Pulseless disease is : (D) Goître (A) PAN 81. The sacro-illac joint transmit from the lower limbs to the : (B) Burger's disease (A) Sacro-illac ligament (D) Raynauds phenomena (C) Lumbosacral ligament (D) Raynauds phenomena (D) Vertebral column of: 82: Lateral boundary of cubital fossa is formed by: (B) Rheumatic fever (A) Biceps brachii (C) Cirrhosis of liver (B) Brachioracialiis (D) Peptic Ulcer (C) Brachioracialiis (D) Peptic Ulcer (D) Pronator teres (A) Bronchial Asthma 83. Skin of nail bed of ring finger is supplied by: (C) Lipoma (A) Lat. half by median, medial half by uhar (D) Bronchitis (B) Medial half by median, lat. half | () | A) Seeding of body cavitied | | 80. | Codman Triangle is a feature of |
| (C)Haematogenous(B)Osteochondroma(D)Lymphatic pathway(C)Chondrosarcoma76.Pulseless disease is :(D)Goitre(A)PAN81.The sacro-iliac joint transmit from the lower limbs to the :(B)Burger's disease(A)Sacro-iliac ligament(C)Takayasus disease(A)Sacro-iliac ligament(D)Raynauds phenomena(C)Lumbosacral ligament77.Anitschkow cells are pathognomonic of:(D)Vertebral columnof:82:Lateral boundary of cubital fossa is formed by:(B)Rheumatic fever(B)Brachioradialis(C)Cirrhosis of liver(B)Brachioradialis(D)Peptic Ulcer(C)Brachioradialis(D)Peptic Ulcer(D)Pronator teres(A)Bronchial Asthma83.Skin of nail bed of ring finger is supplied by:(A)Bronchitis(A)Lat. half by median, medial half by uinar(D)Bronchitis(B)Medial half by median, lat. half by uinar(B)Mallory bodies are seen in : (A)Alcoholic Cirrhosis(C) | (i | 3) Implantation | | ••• | 5 |
| (D)Lymphatic pathway(C)Chondrosarcoma76.Pulseless disease is : (A)(D)Goitre(A)PAN81.The sacro-iliac joint transmit from the lower limbs to the : (A)Sacro-iliac ligament(B)Burger's disease (C)Takayasus disease (D)(A)Sacro-iliac ligament(D)Raynauds phenomena(C)Lumbosacral ligament77.Anitschkow cells are pathognomonic of: (A)(D)Vertebral column61.Rheumatoid arthritis formed by: (B)Rheumatoid arthritis formed by: (B)Brachioradialis (D)(D)Peptic Ulcer(C)Brachioradialis supplied by: (D)Pronator teres78.Reid Index is related to : (A)Bronchial Asthma (D)83.Skin of nail bed of ring finger is supplied by: (B)(B)Lipoma (D)Eipoma (D)C)Lat half by median, medial half by ulnar (B)79.Mallory bodies are seen in : (A)(C)Whole by median nerve | (0 | C) Haematogenous | • | | |
| (A) PAN (B) Burger's disease (C) Takayasus disease (D) Raynauds phenomena 77. Anitschkow cells are pathognomonic of: (A) Rheumatoid arthritis (B) Rheumatic fever (C) Cirrhosis of liver (D) Peptic Ulcer 78. Reid Index is related to : (A) Bronchial Astrma (B) Lipoma (C) Leukemia (D) Bronchitis 79. Mallory bodies are seen in : (A) Alcoholic Cirrhosis | ([| D) Lymphatic pathway | | | |
| (A) PAN (B) Burger's disease (C) Takayasus disease (D) Raynauds phenomena 77. Anitschkow cells are pathognomonic of: (A) Rheumatoid arthritis (B) Rheumatic fever (C) Cirrhosis of liver (D) Peptic Ulcer 78. Reid Index is related to : (A) Bronchial Asthma (B) Lipoma (C) Leukemia (D) Bronchitis 79. Mallory bodies are seen in : (A) Alcoholic Cirrhosis 81. The sacro-iliac joint transmit from the lower limbs to the : (A) Sacro-iliac ligament (B) Pubic symphysis (C) Lumbosacral ligament (D) Vertebral column 82. Lateral boundary of cubital fossa is formed by: (A) Biceps brachii (B) Brachioradialis (C) Brachialis (D) Pronator teres 83. Skin of nail bed of ring finger is supplied by: (A) Lat. half by median, medial half by unar (B) Medial half by median, lat. half by median | 76. P | ulseless disease is : | | | (D) Goitre |
| (B) Burger's disease (C) Takayasus disease (D) Raynauds phenomena 77. Anitschkow cells are pathognomonic of: (A) Rheumatoid arthritis (B) Rheumatic fever (C) Cirrhosis of liver (D) Peptic Ulcer 78. Reid Index is related to : (A) Bronchial Asthma (B) Lipoma (C) Leukemia (D) Bronchitis 79. Mallory bodies are seen in : (A) Alcoholic Cirrhosis | | | | 81 | The sacro-iliac joint transmit from the |
| (C) Takayasus disease (D) Raynauds phenomena (A) Sacro-Iliac ligament (B) Pubic symphysis (C) Lumbosacral ligament (D) Vertebral column (D) Vertebral column (D) Reumatoid arthritis (A) Rheumatic fever (B) Rheumatic fever (C) Cirrhosis of liver (D) Peptic Ulcer (C) Cirrhosis of liver (D) Peptic Ulcer (C) Brachialis (C) Leukemia (D) Bronchitis (A) Alcoholic Cirrhosis (C) Whole by median nerve | | | | | lower limbs to the : |
| (D) Raynauds phenomena (D) Raynauds phenomena (E) Pubic symphysis (C) Lumbosacral ligament (D) Vertebral column (D) Vertebral column 82: Lateral boundary of cubital fossa is formed by: (A) Rheumatic fever (C) Cirrhosis of liver (D) Peptic Ulcer (C) Brachioradialis (D) Peptic Ulcer (C) Brachialis (D) Pronator teres (A) Bronchial Asthma (B) Lipoma (C) Leukemia (D) Bronchitis (A) Alcoholic Cirrhosis (C) Whole by median nerve | | · - | | | (A) Sacro-iliac ligament |
| (C) Lumbosacral ligament (D) Vertebral column (E) Rheumatoid arthritis (A) Rheumatic fever (B) Rheumatic fever (C) Cirrhosis of liver (D) Peptic Ulcer (C) Brachialis (D) Pronator teres (E) Lipoma (C) Leukemia (D) Bronchitis (E) Bronchitis (E) Lipoma (C) Leukemia (D) Bronchitis (E) Mallory bodies are seen in : (A) Alcoholic Cirrhosis (C) Unote the second of the | | | | | (B) Pubic symphysis |
| of:(D) Vertebral courningof:(E) Rheumatoid arthritis(A) Rheumatoid arthritisformed by:(B) Rheumatic fever(A) Biceps brachii(C) Cirrhosis of liver(B) Brachioradialis(D) Peptic Ulcer(C) Brachialis78. Reid Index is related to :(D) Pronator teres(A) Bronchial Asthma83. Skin of nail bed of ring finger is supplied by:(B) Lipoma(A) Lat. half by median, medial half by ulnar(D) Bronchitis(B) Medial half by median, lat. half by median79. Mallory bodies are seen in :(C) Whole by median nerve | | | | - | (C) Lumbosacral ligament |
| 82Lateral boundary of cubital fossa is formed by:(A)Rheumatoid arthritis(B)Rheumatic fever(C)Cirrhosis of liver(D)Peptic Ulcer(D)Peptic Ulcer(C)Bronchial Asthma(B)Lipoma(C)Leukemia(D)Bronchitis(C)Leukemia(D)Bronchitis(C)Leukemia(D)Bronchitis(C)Leukemia(D)Bronchitis(C)Latenal boundary of cubital fossa is formed by:(A)Alcoholic Cirrhosis(A)Alcoholic Cirrhosis(C)Whole by median nerve | ··· | · · · · · · · · · · · · · · · · · · · | iC | | (D) Vertebral column |
| (A) Rheumatoid arthritis (B) Rheumatic fever (C) Cirrhosis of liver (D) Peptic Ulcer (C) Brachialis (C) Brachialis (C) Brachialis (D) Pronator teres (A) Bronchial Asthma (B) Lipoma (C) Leukemia (D) Bronchitis (A) Alcoholic Cirrhosis (C) Whole by median nerve | 0 | f: | | 82. | Lateral boundary of cubital fossa is |
| (C) Cirrhosis of liver (D) Peptic Ulcer 78. Reid Index is related to : (A) Bronchial Asthma (B) Brachioradialis (C) Brachialis (D) Pronator teres 83. Skin of nail bed of ring finger is supplied by: (A) Leukemia (D) Bronchitis 79. Mallory bodies are seen in : (A) Alcoholic Cirrhosis (C) Urbox (C) Whole by median nerve | . (/ |) Rheumatoid arthritis | | | formed by: |
| (D) Peptic Ulcer (C) Brachialis (C) Brachialis (D) Pronator teres (A) Bronchial Asthma (B) Lipoma (C) Leukemia (D) Bronchitis (D) Bronchitis (A) Alcoholic Cirrhosis (B) C) Brachialis (C) Brachialis (D) Pronator teres (C) Leukemia (D) Bronchitis (C) Medial half by median, lat. half (D) Bronchitis (C) Whole by median nerve | , (E | B) Rheumatic fever | | | (A) Biceps brachii |
| 78. Reid Index is related to : (A) Bronchial Asthma (B) Lipoma (C) Leukemia (D) Bronchitis 79. Mallory bodies are seen in : (A) Alcoholic Cirrhosis (D) Pronator teres 83. Skin of nail bed of ring finger is supplied by: (A) Lat. half by median, medial half by ulnar (B) Medial half by median, lat. half by median (C) Whole by median nerve | (0 | Cirrhosis of liver | | | (B) Brachioradialis |
| (A) Bronchial Asthma (B) Lipoma (C) Leukemia (D) Bronchitis 79. Mallory bodies are seen in : (A) Alcoholic Cirrhosis 83. Skin of nail bed of ring finger is supplied by: (A) Lat. half by median, medial half by ulnar (B) Medial half by median, lat. half by median (C) Whole by median nerve | ([|) Peptic Ulcer | • | | (C) Brachialis |
| (B) LipOma (C) Leukemia (D) Bronchitis (A) Lat. half by median, medial half by ulnar (B) Medial half by median, lat. half (B) Medial half by median, lat. half (C) Whole by median nerve | 78. R | eid Index is related to : | | | (D) Pronator teres |
| (B) Eiponna (C) Leukemia (D) Bronchitis (A) Lat. half by median, medial half by ulnar (B) Medial half by median, lat. half (B) Medial half by median, lat. half (C) Whole by median nerve | (A |) Bronchial Asthma | | 83. | Skin of nail bed of ring finger is |
| (C) Leukemia (D) Bronchitis (A) Lat. half by median, medial half by ulnar (B) Medial half by median, lat. half (B) Medial half by median, lat. half (C) Whole by median nerve | (É |) Lipoma | | | supplied by: |
| (D) Bronchitisby ulnar(B) Medial half by median, lat. half79. Mallory bodies are seen in :(A) Alcoholic Cirrhosis(C) Whole by median nerve | (C |) Leukemia | | | (A) Lat. half by median, medial half |
| (B) Medial half by median, lat. half (A) Alcoholic Cirrhosis (B) Medial half by median, lat. half (C) Whole by median nerve | | | | | |
| (A) Alcoholic Cirrhosis (C) Whole by median nerve | | | | | • |
| | | - | | | |
| | | | | | |
| CC- 1A/14 (12) Contd, | | | | | |

Which of the following is not present 84. in the bed of stomach?

小,一个小学的新闻和新闻。 2013

- (A) Splenic artery
- **(B)** Transverse mesocolon
- Fourth part of duodenum (C)
- Transverse colon (D)
- Normal position of uterus is : 85.
 - Anteverted and Anteflexed (A)
 - **Retroverted and Retroflexed (B)**
 - Anteverted and Retroflexed (C)
 - Retroverted and Anteflexed (D)
- 86. There is one sesamoid bone in the tendon of peroneus longus. It articulates with the :
 - (A) Talus
 - Calcaneus **(B)**
 - (C) Navicular
 - Cuboid (D)
- 87. Middle cerebellar peduncle is formed by:
 - (A) Vestibulocerebellar fibers
 - Parolivocerebellar fibers **(B)**
 - Pontocerebellar fibers (C)
 - Posterior spinocerebellar (D) fibers
- CC 1A/14

- The pulmonary veins are four in 88. number, two on each side and they carry:
 - (A) Carbondi-oxided blood
 - **(B)** Carbo-mono-oxided blood
 - (C)Carbo-pera-oxided blood
 - Oxygenated blood (D)
- Which 89. gland is provided secretomotor fibres from inferior salivatory nucleus ?
 - Lacrimal (A)
 - Sublingual (B)
 - (C) Parotid
 - Submandi bular (D)
- 90. All the germ layers, ectoderm, mesoderm and endoderm are present in:
 - (A) Heart
 - **(B)** Cornea
 - Tympanic membrane (C)
 - Urachus. (D)
- 91. The total body water can be measuredusing :
 - (A) Evans Blue
 - Tritium oxide **(B)**
 - Sucrose (C)
 - Sodium thiosulphate (D)

(Turn over)

(13)

- Lymphocytes are concerned with: 92
 - Cell mediated immunity (A)
 - **(B)** Humoral immunity
 - Phagocytosis (C)
 - Delayed hypersensitivity (D)
- Pancreatic enzyme secretion : 93.
 - is mainly from the ductules cells (A)
 - Is from the acinal cells of the **(B)** pancreas
 - Is stimulated by pancreatic (C) polypeptide
 - is stimulated by secretin (D)
- Clearance value of a substance that 94. is freely filtered but neither secreted nor reabsorbed by the kidney will equal to the :
 - Filtration fraction (A)
 - Flomerular filtration rate **(B)**
 - Renal plasma flow (C)
 - Urinary excretion rate of the (D) substance
- 95 When the core temperature of the individual rises more than 41°C (106°F) the following changes occur except:
 - Heat production increases (A)
- CC-1A/14

Sweating increases greatly (B)

- Central nervous system may (C) malfunction
- Person (D) may become poikilothermic
- 96. Which of the following hormones impairs hydrolysis of triglycerides into fatty acids ?
 - Cortisol (A)
 - (B) Glucagon
 - (C) Growth hormone
 - Insulin (D)

Ventricular musle is activated by 97. impulses directly from the :

- (A) Purkinje system
- S-A Node (B)
- Bundle of His (C)
- A-V node (D)
- Histotoxic hypoxia is seen in : 98.
 - Carbon monoxide poisoning (A)
 - Cyanide poisoning **(B)**
 - Shock (C)
 - Anaemias (D)

Contd.

(14)

99. Intestinal absorption of calcium occurs by:

100. In human being, the primary auditory cortex is located in :

- (A) Active uptake (A) Area44
- (B) Simple diffusion
- (C) Facilitated diffusion
- (D) Endocytosis

- (B) Area 41
- (C) Area 43
- (D) Area 22

(Turn over)

4

SPACE FOR ROUGH WORK

1 1 1 1 1

©C - 1A/14 (300)

(16)

Homoeopathic Medical Officer (Paper--I)