

## GGSIPIU Mathematics 2005

1. The equation of the plane through the intersection of the planes  $x+y+z = 1$  and  $2x+3y-z+4 = 0$  and parallel to x-axis is :

- a  $y - 3z + 6 = 0$       b  $3y - z + 6 = 0$   
c  $y + 3z + 6 = 0$       (d)  $3y - 2z + 6 = 0$

2. The distance of the point 3,8,2 from the line  $\frac{x-1}{2} = \frac{y-3}{4} = \frac{z-2}{3}$  measured parallel to the plane  $3x+2y-2z+15 = 0$  is :

- a 2                      b 3  
c 6                      d  $\frac{19}{2}$

3. Let 3,4, -1 and -1,2,3 are the end points of a diameter of sphere. Then the radius of the sphere is equal to :

- a 1      b 2      c 3      d 9

4. If A,B,C,D are the points 2,3, -1,3,5, -3,1,2,3,3,5,7 respectively, Then the angle between AB and CD is :

- a  $\frac{\pi}{2}$                       b  $\frac{\pi}{3}$   
c)  $\frac{\pi}{4}$                       d  $\frac{\pi}{6}$

5. If  $u = \log\left(\frac{x^2+y^2}{x+y}\right)$ , then the value of  $x\frac{\partial u}{\partial x} + y\frac{\partial u}{\partial y}$  is :

- a -1                      b 0  
c 1                      d 2

6. A five digits number is formed by writing the digits 1,2,3,4,5 in a random order without repetitions. Then the probability that the number is divisible by 4, is :

- a  $\frac{3}{5}$                       b  $\frac{18}{5}$   
c  $\frac{1}{5}$                       d  $\frac{6}{5}$

7. Two persons A and B takes turns In throwing a pair of dice. The first person to throw 9 from both dice will be awarded the price. If A throws first, then the probability that B wins the game , is :

- a  $\frac{9}{17}$                       b  $\frac{8}{17}$

c 8/9      d 1/9

8. The probability that in year of the 22<sup>nd</sup> century chosen at random, then there will be 53 Sundays, is :

a 3/28      b 2/28

c 7/28      d 5/28

9. The standard deviation of a variable x is 10. Then the standard deviation of 50+5x is :

a 50      b 550

c 10      d 0.98

10. The octal equivalent of the decimal number 0.3125 is :

a 0.24      b 0.42

c 0.39      d 0.98

11. The hexadecimal equivalent of the binary number 111100001010001 is

a 15C3      b C351

c 3C51      d C315

12. A real value of x will satisfy the equation  $\left(\frac{3-4ix}{3+4ix}\right) = \alpha - i\beta$   $\alpha$  and  $\beta$  are real, if :

a  $\alpha^2 - \beta^2 = -1$       b  $\alpha^2 - \beta^2 = 1$

c  $\alpha^2 + \beta^2 = 1$       d  $\alpha^2 - \beta^2 = 2$

13. If  $\omega$  is a complex cube root of unity, then the value of

$$\frac{p+q\omega+r\omega^2}{r+p\omega+q\omega^2} + \frac{p+q\omega+r\omega^2}{q+r\omega+p\omega^2} \quad p, q, r \in \mathbb{R} \text{ is equal to :}$$

a 0      b 1

c -1      d 2

14. If P, Q, R, S are represented by the complex numbers  $4 + i, 1 + 6i, -4 + 3i, -1 - 2i$  respectively, then PQRS is a :

a rectangle      b square

c rhombus      d parallelogram

15. If n is a positive integer, then  $1+i^n + 1-i^n$  is equal to :

a  $\sqrt{2}^{n-2} \cos\left(\frac{n\pi}{4}\right)$

b  $\sqrt{2}^{n-2} \sin\left(\frac{n\pi}{4}\right)$

c  $\sqrt{2}^{n+2} \cos\left(\frac{n\pi}{4}\right)$

d  $\sqrt{2}^{n+2} \sin\left(\frac{n\pi}{4}\right)$

16. The number of ways in which 9 persons can be divided into three equal groups is :

a 1680    b 840

c 560    d 280

17. A dictionary is printed consisting of 7 letters words only that can be made with a letters of the word CRICKET.If the words are printed are alphabetical order is an ordinary dictionary,then the number of words are before the word CRICKET is :

a 530    b 480

c 531    d 481

18. If the sum of the coefficient in the expansion of  $x+y^n$  is 1024, then the value of the greatest coefficient in the expansion is :

a 356    b 252

c 210    d 120

19. The value of the determinant

$$\begin{vmatrix} 10! & 11! & 12! \\ 11! & 12! & 13! \\ 12! & 13! & 14! \end{vmatrix} \text{ is :}$$

a  $2 \cdot 10! \cdot 11!$     b  $2 \cdot 10! \cdot 13!$

c  $2 \cdot 10! \cdot 11! \cdot 12!$     d  $2 \cdot 11! \cdot 12! \cdot 13!$

20. If A and B are 3x3 matrices such that  $AB = B$  and  $BA = A$ ,than :

a  $A^2 = A$  and  $B^2 \neq B$

b  $A^2 \neq A$  and  $B^2 = B$

c  $A^2 = A$  and  $B^2 = B$

d  $A^2 \neq A$  and  $B^2 \neq B$

21. If the points  $(x_1, y_1)$ ,  $(x_2, y_2)$  and  $(x_3, y_3)$  are collinear, then the rank of the matrix

$$\begin{bmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{bmatrix} \text{ will always be less than :}$$

- a 2      b 3  
c 1      d none of these

22. The system of equations;  $x+y+z = 6, x+2y+3z = 10, x+2y+\lambda z = 6$  has number solution for :

- a  $\lambda = 3, \mu = 10$       b  $\lambda = 3, \mu \neq 10$   
c  $\lambda \neq 3, \mu \neq 10$       d none of these

23. If  $A = \begin{bmatrix} \sin(\theta + \alpha) & \cos(\theta + \alpha) & 1 \\ \sin(\theta + \beta) & \cos(\theta + \beta) & 1 \\ \sin(\theta + \gamma) & \cos(\theta + \gamma) & 1 \end{bmatrix}$  Then :

- a  $A=0$  for all  $\theta$   
b  $A$  is a odd function of  $\theta$   
c  $A=0$  for  $\theta = \alpha + \beta + \gamma$   
d  $A$  is a independent of  $\theta$

24. An investigator interviewed 100 students to determine the performance of three drinks milk, coffee and tea; 20 students take milk and coffee, 30 students take coffee and tea, 25 students take milk and tea, 12 students take milk only, 5 students take coffee only and 8 students take tea only. Then the number of students who did not take any drinks any of three, is :

- a 10                      b 20  
c 25                      d 30

25. Let  $Y = \{1, 2, 3, 4, 5\}$ ,  $A = \{1, 2\}$ ,  $B = \{3, 4, 5\}$  and  $\phi$  denotes null set. If  $A \times B$  denotes Cartesian product of the sets  $A$  and  $B$ , then  $Y \times A \cap Y \times B$  is :

- a  $Y$                       b  $A$   
b  $B$                       d  $\phi$

26. Let  $A = \{2, 3, 4, 5, \dots, 16, 17, 18\}$ . Let  $\approx$  be the equivalence relation on  $A \times A$  Cartesian product of  $A$  and  $A$ , defined by  $a, b \approx c, d$  if  $ad = bc$ , then the number of ordered pairs of the equivalence class of  $(3, 2)$  is :

a 4   b 5   c 6   d 7

27. A question 'who have studied Physics?' was asked to three students A,B and C. The question was answered correctly as it is true that If A studied Physics, then B also studied Physics but it is false statement that if C studied Physics, then B also studied physics. Then physics was studied by :

a both A and B      b only A  
c only B              d only C

28. If a,b be two fixed positive integers such that  $f(a+x) = b + [b^3 + 1 - 3b^2 f x + 3b\{f x\} - \{f(x)^3\}^{1/3}]$  for all real x, then f(x) is a periodic function with period :

a a                      b 2a  
c | b                    d 2b

29. The domain of the function  $f(x) = \log_3 + x \sqrt{x^2 - 1}$  is :

a  $-3, -1 \cup 1, \infty$   
b  $[-3, -1] \cup [1, \infty$   
c  $-3, -2 \cup -2, -1 \cup 1, \infty$   
d  $[-3, -2 \cup -2, -1 \cup 1, \infty$

30. The value of  $\cot 70^\circ + 4 \cos 70^\circ$  is :

a  $1/\sqrt{3}$       b  $\sqrt{3}$   
c  $2\sqrt{3}$       d  $1/2$

31. The equation of  $\sin x + \sin y + \sin z = -3$  for  $0 \leq x \leq 2\pi, 0 \leq y \leq 2\pi, 0 \leq z \leq 2\pi$  has :

a one solution  
b two sets of solution  
c four sets of solution  
d no solution

32. If  $\theta = \sin^{-1} x + \cos^{-1} x - \tan^{-1} x, x \geq 0$  then the smallest interval in which  $\theta$  lies is :

a  $\frac{\pi}{2} \leq \theta \leq \frac{3\pi}{4}$       b  $0 \leq \theta \leq \frac{\pi}{4}$   
c  $-\frac{\pi}{4} \leq \theta \leq 0$       d  $\frac{\pi}{4} \leq \theta \leq \frac{\pi}{2}$

33. Let A,B and C are the angles Of a plain triangle and  $\tan\left(\frac{A}{2}\right) = \frac{1}{3}, \tan\left(\frac{B}{2}\right) = \frac{2}{3}$ . Than  $\tan\left(\frac{C}{2}\right)$  is equal to :

- a 7/9   b 2/9   c 1/3   d 2/3   2/3

34. If  $\alpha, \beta$   $\alpha \neq \beta$  satisfies the question  $a \cos \theta + b \sin \theta = c$ , then the value of  $\tan\left(\frac{\alpha+\beta}{2}\right)$  is :

- a b/a   b c/a   c a/b   d c/b

35. A ray of light passing through the point 1,2 is reflected on the x -axis at a point P and passes through the point 5,3, then the abscissa of a point P is :

- a 3      b 13/3  
c 13/ 5    d 13/4

36. The equation  $4x^2 - 24xy + 11y^2 = 0$  represents :

- a two parallel lines  
b two perpendicular lines  
c two lines through the origin  
d a circle

37. The length of the chord joining the points in which the straight line  $\frac{x}{3} + \frac{y}{4} = 1$  cuts the circle  $x^2 + y^2 = \frac{169}{25}$  is :

- a 1      b 2  
c 4      d 8

38. The normal to the parabola  $y^2 = 8x$  at the point 2,4 meets the parabola against the point :

- a -18,-12    b -18,12  
c 18,12      d 18, -12

39. If a bar of given length moves with its extremities on two fixed straight lines at right angles, then the locus of any point on bar marked on the bar describes a/an :

- a circle      b parabola  
c ellipse     d hyperbola

40. The straight line  $x+y=\sqrt{2}p$  will touch the hyperbola  $4x^2-9y^2=36$  if :

- a  $p^2 = 2$       b  $p^2 = 5$

c  $5p^2 = 2$       d  $2p^2 = 5$

41. The function  $f(x) = \frac{1 - \sin x + \cos x}{1 + \sin x + \cos x}$  is not defined at  $x = \pi$ . The value of  $f(\pi)$ , so that  $f(x)$  is continuous at  $x = \pi$ , is :

- a  $-1/2$       b  $1/2$   
c  $-1$       d  $1$

42. If  $f(x) = \sin^2 x$  and the composite function  $gf(x) = |\sin x|$ , then the function  $g(x)$  is equal to :

- a  $\sqrt{x-1}$       b  $\sqrt{x}$   
c  $\sqrt{x+1}$       d  $-\sqrt{x}$

43. Area of the figure bounded by the curves  $y = |x-1|$  and  $y = 3 - |x|$  is :

- a 1 sq. units      b 2 sq. units  
c 3 sq. units      d 4 sq. units

44. Let  $x = \left[ \frac{a+2b}{a+b} \right]$  and  $y = \frac{a}{b}$ , where  $a$  and  $b$  are positive integers. If  $y^2 > 2$ , then :

- a  $x^2 \leq 2$       b  $x^2 < 2$   
c  $x^2 > 2$       d  $x^2 \geq 2$

45.  $\int_0^1 \tan^{-1} \left( \frac{1}{x-x+1} \right) dx$  is :

- a  $\log 2$       b  $-\log 2$   
c  $\frac{\pi}{2} + \log 2$       d  $\frac{\pi}{2} - \log 2$

46. The curves  $x = \log y + e$  and  $y = \log \left( \frac{1}{x} \right)$  :

- a do not meet  
b meet at one point  
c meet at two points  
d meet at more than two points

47.  $\lim_{x \rightarrow 0} \frac{\cos(\sin x) - 1}{x^2}$  equals :

- a 0      b -1

c  $\frac{1}{2}$  d  $-\frac{1}{2}$

48. Let  $\vec{i}, \vec{j}, \vec{k}$  be three vectors from  $\vec{i} \times \vec{j} \times \vec{k} = \vec{i} \times \vec{j} \times \vec{k}$  it :

a  $\vec{i} \times \vec{j} \times \vec{k} = \vec{i}$  b  $\vec{i} \times \vec{j} \times \vec{k} = \vec{j}$

c  $\vec{i} \times \vec{j} = \vec{i} \times \vec{k}$  d  $\vec{i} \times \vec{j} = \vec{j} \times \vec{k}$

49. If  $\vec{i}, \vec{j}, \vec{k}$  are units vectors and  $|\vec{a}| = a$ , then the value of

$|\vec{i} \times \vec{a}|^2 + |\vec{j} \times \vec{a}|^2 + |\vec{k} \times \vec{a}|^2$  is :

a  $a^2$  b  $3a^2$  c  $2a^2$  d  $4a^2$

50. If the area above the x-axis bounded by the curves  $y = 2^{kx}$  and  $x = 0$  and  $x = 2$  is  $\frac{3}{\log 2}$ , then the value of k is :

a  $\frac{1}{2}$  b 1 c -1 d 2

51. The value of  $\int_a^b \frac{x}{|x|} dx, a < b < 0$  is :

a  $-|a| + |b|$  b  $|b| - |a|$

c  $|a| - |b|$  d  $|a| + |b|$

52. The value of

$\int_{-2}^2 \left[ p \log \left( \frac{1+x}{1-x} \right) + q \log \left( \frac{1-x}{1+x} \right)^{-2} + r \right] dx$  depends on:

a The value of p

b The value of q

c The value of r

d The value of p and q

53. A curve having the condition that the slope of tangent at some point is two times the slope of the straight line joining the same point to the origin of co-ordinates, is a/an :

a circle b ellipse

c parabola d hyperbola

54. If a is an arbitrary constant, then solution of differential equation

$\frac{dy}{dx} + \sqrt{\frac{1-y^2}{1-x^2}} = 0$  is :



- a  $x\sqrt{1-y^2} + y\sqrt{1-x^2} = a$
- b  $y\sqrt{1-y^2} + x\sqrt{1-x^2} = a$
- c  $x\sqrt{1-y^2} - y\sqrt{1-x^2} = a$
- d  $y\sqrt{1-y^2} - x\sqrt{1-x^2} = a$

55. A particle is moving along the curve  $x = at^2 + bt + c$ . If  $a = b^2$ , then the particle would be moving with uniform :

- a rotation      b velocity
- c acceleration      d retardation

56. The solution of the equation  $x^2 \frac{d^2y}{dx^2} = \log x$  when  $x=1, y=0$  and  $\frac{dy}{dx} = -1$  is :

- a  $\frac{1}{2} \log x^2 + \log x$
- b  $\frac{1}{2} (\log x^2 - \log x)$
- c  $-\frac{1}{2} \log x^2 + \log x$
- d  $-\frac{1}{2} \log x^2 - \log x$

57. Let the unit vectors  $\vec{i}$  and  $\vec{j}$  be perpendicular to each other and the unit vector  $\vec{k}$  be inclined at an angle  $\theta$  to both  $\vec{i}$  and  $\vec{j}$ . If  $\vec{k} = \alpha \vec{i} + \beta \vec{j} + \gamma \vec{k}$ , where  $\alpha, \beta, \gamma$  are scalars, then :

- a  $\alpha = \cot \theta, \beta = \sin \theta, \gamma^2 = \cos 2\theta$
- b  $\alpha = \cos \theta, \beta = \cos \theta, \gamma^2 = \cos 2\theta$
- c  $\alpha = \cos \theta, \beta = \sin \theta, \gamma^2 = \cos 2\theta$
- d  $\alpha = \sin \theta, \beta = \cos \theta, \gamma^2 = -\cos 2\theta$

58. If  $y = \frac{1}{\sqrt{a^2-b^2}} \cos^{-1} \left[ \frac{a \cos(x-a)+b}{\rho} \right]$  where  $\theta = a+b \cos x - a$ , then  $\frac{dy}{dx}$  is equal to :

- a  $1/\theta$       b  $2/\theta$
- c  $1/\theta^2$       d  $2/\theta^2$

59. Let K be a set of real number and  $f:K \rightarrow R$  such that for all x any y  $|f(x) - f(y)| \leq |x-y|^5$ . If  $f(3) = 7$ , then the value of f(9) is equal to

- a 5      b 7
- c 9      d 11

60. If  $f(x) = \frac{1}{1-x}$  then the derivative of the composite function  $f[f(x)]$  is equal to :

a 0            b  $\frac{1}{2}$

c 1            d 2